



Rte Ancienne Papeterie
PO Box 327,
CH-1723 Marly 1 (Switzerland)

Fax: +41 91 825 76 08
info@swant.com
www.swant.com

Product Description Monoclonal anti GABA 3A12

Product: monoclonal anti-GABA antibody

Code No: mAB 3A12

Lot no: ps2

Form: Lyophilized (no preservatives).

Quantity: 200 µl.

Reconstitution

- add the lyophilized material (approx. 1 mg) to 100 µl PBS or another buffered saline (add in order indicated)
- allow some time for the material to dissolve; mix by swirling gently
- remove particulate matter by centrifugation in a microfuge
- add 100 µl glycerol (final glycerol conc. 50% v/v) and mix well (vortex)

Description

The antibodies have been purified from serum-free culture medium (Tecnomouse bioreactor) by ammoniumsulfate precipitation, dialyzed against water and lyophilized. mAb 3A12 is not suitable for postembedding immunogold procedures at the EM level.

Storage

Store stock-solution in freezer (-20°C, not -70°C). Mix stock solution well before making aliquots. 10x diluted solutions may be kept in the refrigerator for some time (several weeks) but should be made 10 mM with sodium azide. Volumes of working dilutions (high dilutions) should be calculated such the they are used up the same day. Avoid polystyrene tubes.

Dilutions

Appropriate final dilutions should be determined experimentally. They may be in the range of 1:10,000 to 1:20,000 or even higher for floating sections, and around 1:1,000 or higher for mounted semithin sections (postembedding staining). These values are for combination with the PAP procedure. If the DAB reaction product in the PAP procedure is silver enhanced (see: Liu et al., Histochem., 1989, 90: 427-445), dilutions in the range of 1:5,000 to 1:20,000 or higher should be adequate for postembedding staining of semithin sections. Add 10% bovine serum to the diluted antibody.

References

1. Matute C, Streit P, Monoclonal antibodies demonstrating GABA-like immunoreactivity. Histochemistry 1986;86(2):147-57
2. Liu CJ, Grandes P, Matute C, Cuenod M, Streit P: Glutamate-like immunoreactivity revealed in rat olfactory bulb, hippocampus and cerebellum by monoclonal antibody and sensitive staining method. Histochemistry 1989;90(6):427-45