



Rte Ancienne Papeterie
PO Box 327,
CH-1723 Marly 1 (Switzerland)

Fax: +41 91 825 76 08
info@swant.com
www.swant.com

Product Description

Recombinant Calbindin D-28k

mouse, 100 µg

(AgCB100abs)

For immunoblot and absorption experiments

Product: HPLC-purified calbindin D-28k (recombinant, 100 µg) produced in *Escherichia Coli*, after *in vitro* cDNA synthesis.

Use: This protein can be used for immunoblots and absorption experiments. For intracellular injections in neurons we suggest using our other products: AgCB10ic or AgCB100ic.

Lot No.: 13 (20-17)


Form: 100 µg in 250 µl of 20 mM NaHCO₃ with **high CaCl₂** (0.1 mM). Lyophilized.

Reconstitution: with 250 µl of bi-distilled water.

Storage: After reconstitution, freeze in small aliquots (e.g. 1 to 10 µg) and keep at -80°C. Avoid repeated freezing and thawing.

Description: This recombinant protein has been purified by HPLC and migrates as a single band of approximately 28 kDa. The figure shows a gel stained with Coomassie-blue.

SDS-PAGE of HPLC-purified mouse recombinant calbindin D-28k. The arrow marks the full-length native calbindin D-28k (28 kDa).



For adsorption experiments aimed at proofing the specificity of the immunostaining, we suggest the following procedure:

- Dilute 1 µl of one of the Swants antibodies against calbindin D-28k in 5 ml of the usual buffer for immunohistochemistry (final dilution 1:5'000).
- Add 1 µg of the recombinant protein to 1 ml of the diluted antibody solution and mix well.
- Incubate for at least 6 hours in the cold.
- Apply to tissue-sections and incubate for 3 days.
- Complete the immunohistochemical reaction as usual (biotinylated secondary antibody, Streptavidin-Cy2 (or other fluorescent dyes), or peroxidase (followed by DAB-H₂O₂)).

As a result, the immunostaining should be strongly reduced or even completely prevented.